

Table S5. Antimicrobial susceptibility (inhibition zone diameter; mm) of *Paraniabella aurantiaca* CJ426^T

Antibiotics	CJ426 ^T	Susceptible/Resistant*
Amoxicillin (10 µg)	6.0	R
Cephalexin (30 µg)	42.9	S
Chloramphenicol (10 µg)	11.1	R
Ciprofloxacin (5 µg)	32.9	S
Clindamycin (2 µg)	30.3	S
Colistin (10 µg)	6.0	R
Erythromycin (10 µg)	6.0	R
Fosfomycin (200 µg)	6.0	R
Gentamicin (10 µg)	16.9	I
Linezolid (10 µg)	10.5	R
Meropenem (10 µg)	11.8	R
Rifampicin (5 µg)	9.0	R
Streptomycin (5 µg)	6.0	R
Sulfamethoxazole (50 µg)	19.7	S
Tetracycline (30 µg)	11.7	I
Trimethoprim (5 µg)	6.0	R
Tylosin (30 µg)	27.5	S
Vancomycin (5 µg)	6.0	R

*Susceptibility results were recorded as susceptible (S), intermediate (I) or resistant (R) based on breakpoints for *Enterobacteriaceae* and *Staphylococcus* spp. according to the CLSI supplement M100.